

# 1 Interpolation and Curve Fitting

1. For the given data points;

$x$	$y$
1	1.06
2	1.12
3	1.34
5	1.78

- construct the interpolating cubic  $P_3(x) = ax^3 + bx^2 + cx + d$ .

**Hint:** First, write the set of equations then solve it by writing/using a MATLAB program.

- Interpolate for  $x = 4$
- Extrapolate for  $x = 5.5$

Solution:

```
>> A=[1 1 1 1; 8 4 2 1; 27 9 3 1;125 25 5 1]
>> B=[1.06 1.12 1.34 1.78]',
>> X=uptrbk(A,B)
X =
    -0.0200
     0.2000
    -0.4000
     1.2800
>> X'*[27 9 3 1]',
ans =      1.3400
>> X'*A(3,1:4)'
ans =      1.3400
>> X'*[4^3 4^2 4 1]',
ans =      1.6000
>> X'*[(5.5)^3 (5.5)^2 5.5 1]',
ans =      1.8025
```

2. We have given the following MATLAB code to evaluate the Lagrange polynomial  $P(x) = \sum_{k=0}^N y_k \frac{\prod_{j \neq k} (x - x_j)}{\prod_{j \neq k} (x_k - x_j)}$  based on  $N + 1$  points  $(x_k, y_k)$  for  $k = 0, 1, \dots, N$ .

```
function [C,L]=lagran(X,Y)
%Input - X is a vector that contains a list of abscissas
```

```

%           - Y is a vector that contains a list of ordinates
%Output - C is a matrix that contains the coefficents of
%           the Lagrange interpolatory polynomial
%           - L is a matrix that contains the Lagrange coefficient polynomials

w=length(X);
n=w-1;
L=zeros(w,w);
%Form the Lagrange coefficient polynomials
for k=1:n+1
    V=1;
    for j=1:n+1
        if k~=j
            V=conv(V,poly(X(j)))/(X(k)-X(j));
        end
    end
    L(k,:)=V;
end
%Determine the coefficients of the Lagrange interpolator polynomial
C=Y*L;

```

where

- The *poly* command creates a vector whose entries are the coefficients of a polynomial with specified roots.

```

>>P=poly(2)
>> 1 -2
>>Q=poly(3)
>> 1 -3

```

- The *conv* command produces a vector whose entries are the coefficients of a polynomial that is the product of two other polynomials.

```

>>conv(P,Q)
>> 1 -5 6 %Thus the product of P(x) and Q(x) is x^2-5x+6

```

Study this MATLAB code and then use the data set in the previous item to

- interpolate for  $x = 4$
- extrapolate for  $x = 5.5$

Solution:

save with the name *lagran.m*. Then;

```
>> X=[1 2 3 5]
>> Y=[1.06 1.12 1.34 1.78]
>> [C,L]=lagran(X,Y)
C =      -0.0200      0.2000     -0.4000      1.2800
L =
   -0.1250      1.2500     -3.8750      3.7500
    0.3333     -3.0000      7.6667     -5.0000
   -0.2500      2.0000     -4.2500      2.5000
    0.0417     -0.2500      0.4583     -0.2500
>> C*A(3,1:4)'
ans =      1.3400
>> C*[4^3 4^2 4 1]',
ans =      1.6000
>> C*[(5.5)^3 (5.5)^2 5.5 1]',
ans =      1.8025
```

3. We have given the following MATLAB code to construct and evaluate divided-difference table for the (Newton) polynomial of  $\text{degree} \leq N$  that passes through  $(x_k, y_k)$  for  $k = 0, 1, \dots, N$ :

$$P(x) = d_{0,0} + d_{1,1}(x-x_0) + d_{2,2}(x-x_0)(x-x_1) + \dots + d_{N,N}(x-x_0)(x-x_1)\dots(x-x_{N-1})$$

where

$$d_{k,0} = y_k \text{ and } d_{k,j} = \frac{d_{k,j-1} - d_{k-1,j-1}}{x_k - x_{k-j}}$$

```
function [C,D]=newpoly(X,Y)
%Input - X is a vector that contains a list of abscissas
%       - Y is a vector that contains a list of ordinates
%Output - C is a vector that contains the coefficients
%       of the Newton interpolatory polynomial
%       - D is the divided difference table

n=length(X);
D=zeros(n,n);
D(:,1)=Y';

%Use the formula above to form the divided difference table
for j=2:n
    for k=j:n
        D(k,j)=(D(k,j-1)-D(k-1,j-1))/(X(k)-X(k-j+1));
    end
end
```

```

    end
end

%Determine the coefficients of the Newton interpolatory polynomial
C=D(n,n);
for k=(n-1):-1:1
    C=conv(C,poly(X(k)));
    m=length(C);
    C(m)=C(m)+D(k,k);
end

```

Study this MATLAB code and then use the data set in the first item to

- construct the divided-difference table by *hand*
- run the MATLAB code and compare with your table
- interpolate for  $x = 4$
- extrapolate for  $x = 5.5$

Solution:

save with the name *newpoly.m*. Then;

```

>> X=[1 2 3 5]
>> Y=[1.06 1.12 1.34 1.78]
>> [C,D]=newpoly(X,Y)
C =      -0.0200      0.2000     -0.4000      1.2800
D =
      1.0600          0          0          0
      1.1200      0.0600          0          0
      1.3400      0.2200      0.0800          0
      1.7800      0.2200          0     -0.0200
>> C*A(4,1:4)'
ans =      1.7800
>> C*[4^3 4^2 4 1]',
ans =      1.6000
>> C*[(5.5)^3 (5.5)^2 5.5 1]',
ans =      1.8025

```